GENERAL INDEX TO VOLUME VII

New scientific names of plants and the final members of new combinations are printed in **bold face** type; synonyms and page numbers having reference to figures and plates, in *italic;* and previously published scientific names and all other matter, in ordinary type.

A

abietina (*Hymenochaete*), 186
abietina (*Thelephora*), 186
abietinum (*Stereum*), 186, *246*
abnormis (*Hymenochaete*), 186
acerinum (*Stereum*), 236
acerinum var. *nivosum* (*Stereum*), 236
Acid, in relation to growth of bacteria, 251, 258, 281, 287
Actinomyces chromogenus, 252
aculeata (Thelephora), 232
affinis (Thelephora), 96
affinis (*Thelephora*), 96
albobadia (*Thelephora*), 216
albobadium (Stereum), 216, *248*
albo-marginata (*Peniophora*), 216
albo-marginata (*Thelephora*), 216
Aleurodiscus subcruentatus, 237
Alkali, in relation to growth of bacteria, 251, 258, 281, 287
Allen, E. R. On carbohydrate consumption by Azotobacter chroococcum, 75
ambigua (Hymenochaete), 200
ambiguum (Stereum), 190, *246*
ambiguus (*Trichocarpus*), 200
ambiguus (*Xerocarpus*), 200
anastomosans (Stereum), 115
anastomosans (*Thelephora*), 115
annosum (Stereum), 226
Apparatus, Millikan dew point, 53; rotating, 76
arcticum (*Stereum*), 155
arenicolum (Stereum), 232
areolatum (Stereum), 202
aschistum (*Corticium*), 203
Atkinsonii (*Peniophora*), 200
aurantiaca (*Podoscypha*), 96
aurantiaca (*Thelephora*), 75
aurantiacum (Stereum), 95, *240*
Auricularia reflexa, 150
australe (Stereum), 141, *244*
Autotoxin of bacteria, 250, 258, 265, 281, 287
Azotobacter chroococcum, carbohydrate consumption by, 75

B

Bacillus aerogenes, 252, 255, in plain and dextrose bouillon, growth and hydrogen-ion concentration of, 275; coli, 250, in plain and dextrose bouillon, growth and hydrogen-ion concentration of, 255, 258, 281, 287; diphtheriae, 253; paratyphosis, 250; radicicola, 252; subtilis, 252; typhosus, 251
Bacterial inhibition by metabolic products, 249
Balanced solutions, insoluble salts in, 307
balsameum (*Stereum*), 145
balsameum f. *reflexum* (*Stereum*), 145
Beet decoction, titration curve of, 301
Berkeleyi (*Peniophora*), 203
Bertolonii (*Stereum*), 169
bicolor (*Stereum*), 117
bicolor (*Thelephora*), 117
bizonatum (*Stereum*), 216
Bresadolina, 81; *pallida*, 104
Burtianum (Stereum), 93, *240*

C

caespitosum (Stereum), 116, *244*
calyculus (*Stereum*), 236
candidum (*Stereum*), 236
caperata (*Thelephora*), 87
caperatum (Stereum), 87, *240*
Carbohydrate consumption, by Azotobacter, 75
carolinense (*Stereum*), 236
Ceratostomella coerulea, 71
Chailletii (Stereum), 200, *248*
Chailletii (*Thelephora*), 200
Chambers, W. H. Studies in the physiology of the fungi. XI. Bacterial inhibition by metabolic products, 249
cinerascens (Stereum), 203, *248*
cinerascens (*Thelephora*), 203
cinerescens (*Peniophora*), 203
Cladoderris infundibuliformis, 88
coffearum (*Stereum*), 216
coffeatum (*Stereum*), 117
complicatum (*Stereum*), 169
concolor (Stereum), 163
conicum (Stereum), 179, *246*
Coniophora cerebella, 71
Corn, food reserve in, 291; nutrition experiments with, 6, 310
corrugata (Thelephora), 181
Corticium aschistum, 203; *ephebium*, 204; *lilacino-fuscum*, 229; *Nyssae*, 128;

ochroleucum, 235; *pezizoideum*, 121; *siparium*, 128; *subrepandum*, 229; *subzonatum*, 150
Cotyledons, nutritive value of the food reserve in, 291
craspedia (*Thelephora*), 112
craspedium (Stereum), 113, *242*
crassa (*Hymenochaete*), 192
crassa (*Thelephora*), 192
crassum (Stereum), 180
Craterella pallida, 104
crateriformis (*Hymenochaete*), 89
cristatum (Stereum), 103
cristulatum (*Stereum*), 136
Crone's solution, 3, 309
Cryptochaete rufa, 121
Culture media, liquid, titration curves of, 299
cuneatum (Stereum), 233
cupulatum (Stereum), 179, 233
cyphelloides (Stereum), 112, *242*
Czapek's solution, titration curve of, 301

D

decolorans (*Podoscypha*), 107
decolorans (Stereum), 107, *242*
decolorans (*Thelephora*), 107
diaphana (*Thelephora*), 98
diaphanum (Stereum), 97, *240*
dissita (*Peniophora*), 203
dissitum (*Stereum*), 203
Duggar, B. M. Hydrogen-ion concentration and the composition of nutrient solutions in relation to the growth of seed plants, 1; The nutritive value of the food reserve in cotyledons, 291; The use of "insoluble"salts in balanced solutions for seed plants, 307
durum (Stereum), 226, *248*

E

Earlei (Stereum), 199, *248*
elegans (Stereum), 93, 105, *242*
elegans (*Thelephora*), 105, 107
Ellisii (*Peniophora*), 222
ephebia (*Peniophora*), 204
ephebium (*Corticium*), 204
erumpens (Stereum), 209, *248*
exigua (*Thelephora*), 99
exiguum (Stereum), 99, *240*

F

fasciatum (Stereum), 155, *246*
ferreum (Stereum), 202, *248*
Ferric citrate, value of, in culture media, 321
Ferric phosphate, soluble, value of, in culture media, 321
fimbriata (*Hymenochaete*), 186
fimbriatum (Stereum), 234
fissum (Stereum), 111, *242*
flabellata (*Podoscypha*), 111
flabellatum (Stereum), 111
Food reserve in cotyledons, nutritive value of, 291
fragile (Stereum), 233
frustulosa (*Thelephora*), 227
frustulosum (Stereum), 227, *248*
fulvo-nitens (*Stereum*), 91
Fungi, studies in the physiology of the, XI, 249
fusca (*Thelephora*), 117
fuscum (Stereum), 117, *244*

G

Galeottii (Stereum), 234
gausapata (*Thelephora*), 136
gausapatum (Stereum), 136, *244*
Geographic distribution of Stereums, 82
glabrescens (Stereum), 110, *242*
glaucescens (*Stereum*), 186
griseum (Stereum), 234
Growth curves of bacteria, 249
Growth of: Azotobacter chroococcum, 75; corn in nutrient solutions, 6, 310; peas in nutrient solutions, 6; wheat in nutrient solutions, 6, 309; seedlings, effects of cotyledons upon, 291
guadelupense (*Stereum*), 236

H

Hartmanni (Stereum), 112, *242*
Hartmanni (*Thelephora*), 112
Haydeni (Stereum), 236
Helvella pannosa, 104; *versicolor*, 167
heterosporum (Stereum), 220, *248*
hirsuta (*Thelephora*), 150
hirsuta var. *ramealis* (*Thelephora*), 169
hirsutum (Stereum), 150, *246*
Huberianum (*Stereum*), 111
Humidity, in relation to moisture imbibition by wood and to spore germination on wood, 51; determination of, 53
Humidors used in wood-decay experiments, description of, 52; view of, 74
Hydrogen-ion concentration, and the composition of nutrient solutions in relation to the growth of seed plants, 1, 327; in relation to growth of bacteria, 252, 256, 258, 281, 287
hydrophorum (Stereum), 89, *240*
Hymenochaete *abietina*, 186; *abnormis*, 186; ambigua, 200; *crassa*, 192; *crateriformis*, 89; *fimbriata*, 186; *Kalchbrenneri*, 192; *multispinulosa*, 192; pallida, 196; *paupercula*, 216; *purpurea*, 192;

rugispora, 188; *scabriseta*, 192; *umbrina*, 192; *vinosa*, 192
Hypocrea Richardsonii, 121

I

illudens (Stereum), 225
Imbibition by wood, relation of, to humidity, 51
infundibuliformis (Cladoderris), 88
insigne (Stereum), 225, *248*
insolitum (*Stereum*), 237
"Insoluble" salts in balanced solutions, 307
intermedia (*Peniophora*), 192

K

Kalchbrenneri (*Hymenochaete*), 192
Karrer, J. L., and R. W. Webb. Titration curves of certain liquid culture media, 299
Kneiffia purpurea, 192

L

lamellata (*Thelephora*), 88
lamellatum (Stereum), 88
Lenzites saepiaria, humidity and spore germination of, 51
Leveillianum (*Stereum*), 237
lilacino-fuscum (*Corticium*), 229
lilacino-fuscum (*Stereum*), 229
Lloydella, 81; *occidentalis*, 204; *scabriseta*, 192; *striata*, 186
lobata (*Thelephora*), 163, 169
lobatum (Stereum), 163, *246*

M

macrorrhiza (*Thelephora*), 93
macrorrhizum (Stereum), 92
magnisporum (Stereum), 207, *248*
Mancianus (*Stereum*), 237
Media, liquid culture, titration curves of, 299
Merulius lacrymans, 71
Metabolic products, bacterial inhibition by, 249
Micheneri (*Stereum*), 128, 237
Mineral nutrient solutions, 1, 307
Moisture content of wood, 54; determination of, 54
molle (*Stereum*), 155
mollis (*Thelephora*), 155
moricola (*Peniophora*), 203
moricola (*Stereum*), 203
multispinulosa (*Hymenochaete*), 192
Murraii (*Thelephora*), 131
Murrayi (Stereum), 131, *244*
mytilina (Thelephora), 141

N

neglecta (*Peniophora*), 204
neglectum (*Stereum*), 204
nicaraguense (*Stereum*), 196
nitidulum (Stereum), 101
Nutrient solutions for seed plants, 1, 307
Nyssae (*Corticium*), 128

O

obscura (Peniophora), 222
obscura (*Thelephora*), 222
occidentale (*Stereum*), 136
occidentalis (*Lloydella*), 204
occidentalis (*Peniophora*), 204
ochracea (*Thelephora*), 150
ochraceo-flava (*Thelephora*), 183
ochraceo-flavum (Stereum), 183, *246*
ochroleucum (*Corticium*), 235
ochroleucum (Stereum), 148, 235
ostrea (*Stereum*), 155
ostrea (*Thelephora*), 155

P

pallida (*Bresadolina*), 104
pallida (*Craterella*), 104
pallida (Hymenochaete), 196
pallida (*Thelephora*), 104
pallidum (Stereum), 104, *242*
pannosa (*Helvella*), 104
pannosa (*Thelephora*), 104
papyrina (*Peniophora*), 196
papyrinum (Stereum), 196, *248*
patelliforme (Stereum), 182, *246*
paupercula (*Hymenochaete*), 216
paupercula (*Peniophora*), 216
Peas, Canada field, food reserve of, 291; nutrition experiments with, 6
Peniophora albo-marginata, 216; *Atkinsonii*, 200; *Berkeleyi*, 203; *cinerescens*, 203; *dissita*, 203; *Ellisii*, 222; *ephebia*, 204; *intermedia*, 192; *moricola*, 203; *neglecta*, 204; obscura, 222; *occidentalis*, 204; *papyrina*, 196; *paupercula*, 216; *Schweinitzii*, 203
Peptone solution, titration curve of, 301
perdix (*Thelephora*), 227
pergamenum (Stereum), 101, *240*
petalodes (Stereum), 114, *242*
pezizoidea (*Tubercularia*), 121
pezizoideum (*Corticium*), 121
Pfeffer's solution, titration curve of, 301
Pini (*Sterellum*), 123
Pini (Stereum), 123, *244*
Pini (*Thelephora*), 123
Pinus echinata, imbibition of wood of, 51; palustris, imbibition of wood of, 51
Pneumococcus, 253

Podoscypha, 81; *aurantiaca*, 96; *decolorans*, 107; *flabellata*, 111; *radicans*, 108
populneum (Stereum), 237
prolifera (*Thelephora*), 115
proliferum (Stereum), 115, *244*
pruinatum (Stereum), 237
pubescens (Stereum), 178, *246*
pulverulentum (*Stereum*), 131
purpurascens (*Stereum*), 204
purpurea (*Hymenochaete*), 192
purpurea (*Kneiffia*), 192
purpurea (*Thelephora*), 124
purpureum (Stereum), 124, *244*
pusiolum (Stereum), 109, *242*

Q

quisquiliare (Stereum), 95, *240*
quisquiliaris (*Thelephora*), 95

R

radians (*Stereum*), 167
radiatum (Stereum), 181, *246*
radiatum var. *reflexum* (*Stereum*), 181
radicans (*Podoscypha*), 108
radicans (Stereum), 108, *242*
radicans (*Thelephora*), 108
rameale (Stereum), 169, *246*
Ravenelii (Stereum), 90, *240*
reflexa (*Auricularia*), 150
Resin content of wood, determination of, 54; in relation to moisture absorption, 66
Richards' solution, titration curve of, 301
Richardsonii (*Hypocrea*), 121
rigens (*Stereum*), 145
rivulorum (Stereum), 94
roseo-carnea (*Thelephora*), 229
roseo-carneum (Stereum), 229, *248*
rufa (*Cryptochaete*), 121
rufa (*Thelephora*), 121
rufum (*Stereum*), 120, *244*
rugispora (*Hymenochaete*), 188
rugisporum (Stereum), 188, *248*
rugosa (*Thelephora*), 143
rugosiusculum (Stereum), 127, *244*
rugosum (Stereum), 142, *244*

S

Salt requirements of seed plants, 1
sanguinolenta (*Thelephora*), 145
sanguinolentum (Stereum), 144, *246*
saxitas (Stereum), 134, *244*
scabriseta (*Hymenochaete*), 192
scabriseta (*Lloydella*), 192
Schweinitzii (*Peniophora*), 203
scriblitum (*Stereum*), 237
Seed plants, hydrogen ion concentration and the composition of nutrient solutions in, 1; "insoluble" salts in balanced solutions for, 307
sendaiense (*Stereum*), 229
sepium (Stereum), 215, *248*
seriatum (*Stereum*), 237
sericella (*Thelephora*), 96
sericeum (Stereum), 175, *246*
Shive's solution, 3
siparium (*Corticium*), 128
Solutions, balanced, "insoluble" salts in, for seed plants, 307
Sowerbeyi (*Stereum*), 104
Sowerbeyi (*Thelephora*), 104
spadicea (*Thelephora*), 136
spadiceum (*Stereum*), 136
spadiceum var. *plicatum* (*Stereum*), 136
spectabilis (*Thelephora*), 96
spongiosum (*Stereum*), 237
Spore germination on wood, 68
Sprucei (*Stereum*), 163
spumeum (Stereum), 208, 236, *248*
Sterellum Pini, 123
Stereum, 81; abietinum, 186, *246; aceri-inum*, 236, var. *nivosum*, 236; albobadium, 216, *248;* ambiguum, 190, *246;* anastomosans, 115; annosum, 226; *arcticum*, 155; arenicolum, 232; areolatum, 202; aurantiacum, 95, *240;* australe, 141, *244; balsameum*, 145, f. *reflexum*, 145; *Bertolonii*, 169; *bicolor*, 117; *bizonatum*, 216; Burtianum, 93, *240;* **caespitosum**, 116, *244; calyculus*, 236; *candidum*, 236; caperatum, 87, *240; carolinense*, 236; Chailletii, 200, *248;* cinerascens, 203, *248; coffearum*, 216; *coffeatum*, 117; *complicatum*, 169; concolor, 163; **conicum**, 179, *246;* craspedium, 113, *242;* crassum, 180; cristatum, 103; *cristulatum*, 136; cuneatum, 233; cupulatum, 179, 233; cyphelloides, 112, *242;* decolorans, 107, *242;* diaphanum, 97, *240; dissitum*, 203; **durum**, 226, *248;* **Earlei**, 199, *248;* elegans, 93, 105, *242;* **erumpens**, 209, *248;* exiguum, 99, *240;* fasciatum, 155, *246;* ferreum, 202, *248;* fimbriatum, 234; fissum, 111, *242;* flabellatum, 111; fragile, 233; frustulosum, 227, *248; fulvo-nitens*, 91; fuscum, 117, *244;* Galeottii, 234; gausapatum, 136, *244;* glabrescens, 110, *242; glaucescens*, 186; griseum, 234; *guadelupense*, 236; Hartmanni, 112, *242;* Haydeni, 236; **heterosporum**, 220, *248;* hirsutum, 150, *246; Huberianum*, 111; hydrophorum, 89, *240;* illudens, 225; insigne, 225, *248; insolitum*, 237; lamellatum, 88; *Leveillianum*, 237; *lilacino-fuscum*, 229; lobatum, 163, *246;* macrorrhizum, 92; **magnisporum**, 207, *248; Mancianus*, 237; *Micheneri*, 128, 237; *molle*, 155; *moricola*, 203; Murrayi, 131, *244; neg-*

lectum, 204; *nicaraguense*, 196; nitidulum, 101; *occidentale*, 136; ochraceoflavum, 185, *248;* ochroleucum, 148, 235; *ostrea*, 155; pallidum, 104, *242;* papyrinum, 191, *248;* **pateliiforme**, 182, *246;* pergamenum, 101, *240;* petalodes, 114, *242;* Pini, 123, *244;* populneum, 237; proliferum, 115, *244;* pruinatum, 237; **pubescens**, 178, *246;* *pulverulentum*, 131; *purpurascens*, 204; purpureum, 124, *244;* pusiolum, 109, *242;* quisquiliare, 95, *240;* radians, 167; radiatum, 181, *246;* *radiatum* var. *reflexum*, 181; radicans, 108, *242;* rameale, 169, *246;* Ravenelii, 90, *240;* *rigens*, 145; rivulorum, 94; roseocarneum, 229, *248;* rufum, 120, *244;* rugisporum, 188, *248;* rugosiusculum, 127, *244;* rugosum, 142, *244;* sanguinolentum, 144, *246;* **saxitas**, 134, *244;* *scriblitum*, 237; *sendaiense*, 229; **sepium**, 215, *248;* *seriatum*, 237; sericeum, 175, *246;* *Sowerbeyi*, 104; *spadiceum*, 136, var. *plicatum*, 136; *spongiosum*, 237; *Sprucei*, 163; **spumeum**, 208, 236, *248;* *strumosum*, 237; styracifluum, 135, *244;* *subcruentatum*, 237; subpileatum, 213, *248;* sulcatum, 211, *248;* *Sullivantii*, 98; sulphuratum, 148, *246;* surinamense, 91, *240;* tenerrimum, 100, *240;* triste, 238; *tuberculosum*, 131; umbrinum, 191, *248;* undulatum, 100; unicum, 236; *variicolor*, 150; versicolor, 166, *246;* versiforme, 222, *248;* vibrans, 179, *246;* *vorticosum*, 124; *Willeyi*, 98; xanthellum, 96

Streptococcus erysipelatus, 252; hemolyticus, 253; lacticus, 252

striata (*Lloydella*), 186

striata (*Thelephora*), 186

striatum (*Stereum*), 186

strumosum (*Stereum*), 237

Studies in the physiology of the fungi. XI, 249

styraciflua (*Thelephora*), 135

styracifluum (Stereum), 135, *244*

subcruentatum (*Stereum*), 237

subcruentatus (Aleurodiscus), 237

subpileatum (Stereum), 213, *248*

subrepandum (*Corticium*), 229

subzonata (*Thelephora*), 150

subzonatum (*Corticium*), 150

sulcatum (Stereum), 211, *248*

Sullivantii (*Stereum*), 98

sulphuratum (Stereum), 148, *246*

surinamense (Stereum), 91, *240*

T

tenerrimum (Stereum), 100, *240*

Thelephora *abietina*, 186; aculeata, 232; affinis, 96; *affinis*, 96; *albobadia*, 216; *albo-marginata*, 216; *anastomosans*, 115; *aurantiaca*, 75; *bicolor*, 117; *caperata*, 87; *Chailletii*, 200; *cinerascens*, 203; *corrugata*, 181; *craspedia*, 112; *crassa*, 192; *decolorans*, 107; *diaphana*, 98; *elegans*, 105, 107; *exigua*, 99; *frustulosa*, 227; *fusca*, 117; *gausapata*, 136; *Hartmanni*, 112; *hirsuta*, 150, var. *ramealis*, 169; *lamellata*, 88; *lobata*, 163, 169; *macrorrhiza*, 93; *mollis*, 155; *Murrayi*, 131; mytilina, 141; *obscura*, 222; *ochracea*, 150; *ochracea-flava*, 183; *ostrea*, 155; *pallida*, 104; *pannosa*, 104; *perdix*, 227; *Pini*, 123; *prolifera*, 115; *purpurea*, 124; *quisquiliaris*, 95; *radicans*, 108; *roseo-carnea*, 229; *rufa*, 121; *rugosa*, 143; *sanguinolenta*, 145; *sericella*, 96; *Sowerbeyi*, 104; *spadicea*, 136; *spectabilis*, 96; *striata*, 186; *styraciflua*, 135; *subzonata*, 150; *versicolor*, 167, var. *fasciata*, 155

Titration curves, culture media, 299

Trichocarpus ambiguus, 200

triste (Stereum), 238

Tuberculariа pezizoidea, 121

tuberculosum (*Stereum*), 131

U

umbrina (*Hymenochaete*), 192

umbrinum (Stereum), 191, *248*

undulatum (Stereum), 100

unicum (Stereum), 236

V

variicolor (*Stereum*), 150

versicolor (*Helvella*), 167

versicolor (Stereum), 166, *246*

versicolor (*Thelephora*), 167

versicolor var. *fasciata* (*Thelephora*), 155

versiforme (Stereum), 222, *248*

vibrans (Stereum), 179, *246*

vinosa (*Hymenochaete*), 192

Vitamine, in relation to cotyledons, 297

vorticosum (*Stereum*), 124

W

Webb, R. W., J. L. Karrer, and. Titration curves in certain liquid culture media, 299

Wheat, nutrition experiments with, 6, 309

Willeyi (*Stereum*), 98

Wood, imbibition in relation to humidity of, 51

Wood-destroying fungi, moisture in relation to, 51

X

xanthellum (Stereum), 96

Xerocarpus ambiguus, 200

Z

Zeller, S. M. Humidity in relation to moisture imbibition by wood and to spore germination on wood, 51